

AD-A194 569


LIMITING DISTRIBUTIONS OF NON-LINEAR VECTOR FUNCTIONS  
OF STATIONARY GAUSS. (U) NORTH CAROLINA UNIV AT CHAPEL  
HILL CENTER FOR STOCHASTIC PROC. H HO ET AL. MAR 88  
TR-228 AFOSR-TR-88-0418 F49620-95-C-0144 F/G 12/3

1/1

UNCLASSIFIED

NL





MICROCOPY RESOLUTION TEST CHART  
NBS 1963-A

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

AD-A194 569

②

## REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED		1b. RESTRICTIVE MARKINGS DTIC ELL-1027	
2a. SECURITY CLASSIFICATION AUTHORITY NA		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for Public Release; Distribution Unlimited	
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE NA		5. MONITORING ORGANIZATION REPORT NUMBER(S) AFOSR-TR. 88-0418	
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Technical Report No. 228		7a. NAME OF MONITORING ORGANIZATION AFOSR/NM	
6a. NAME OF PERFORMING ORGANIZATION University of North Carolina	6b. OFFICE SYMBOL (If applicable)	7b. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB, DC 20332-6448	
8a. NAME OF FUNDING/SPONSORING ORGANIZATION AFOSR	8b. OFFICE SYMBOL (If applicable) NM	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER F49620 85 C 0144	
8c. ADDRESS (City, State and ZIP Code) Bldg. 410 Bolling AFB, DC		10. SOURCE OF FUNDING NOS.	
		PROGRAM ELEMENT NO. 6.1102F	PROJECT NO. 2304
		TASK NO. H5	WORK UNIT NO. Leave blank
11. TITLE (Include Security Classification) Limiting distributions of non-linear vector functions of stationary Gaussian processes			
12. PERSONAL AUTHOR(S) Ho, H.-C. and Sun, T.-C.			
13a. TYPE OF REPORT preprint	13b. TIME COVERED FROM 9/87 TO 8/88	14. DATE OF REPORT (Yr., Mo., Day) Mar. 1988	15. PAGE COUNT 16
16. SUPPLEMENTARY NOTATION N/A			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB. GR.	
XXXXXXX	XXXXXXX	Key words and phrases: central limit theorem, non-central limit theorem, long-range dependence, stationary Gaussian vector processes	
19. ABSTRACT (Continue on reverse if necessary and identify by block number)			
<p><b>Abstract:</b> Given a stationary Gaussian vector process <math>(X_m, Y_m)</math>, <math>m \in Z</math>, and two real functions <math>H(x)</math> and <math>K(x)</math> we define <math>Z_H^n = A_n^{-1} \sum_{m=0}^{n-1} H(X_m)</math> and <math>Z_K^n = B_n^{-1} \sum_{m=0}^{n-1} K(Y_m)</math>, where <math>A_n</math> and <math>B_n</math> are some appropriate constants. The joint limiting distribution of <math>(Z_H^n, Z_K^n)</math> is investigated. It is shown that <math>Z_H^n</math> and <math>Z_K^n</math> are asymptotically independent when one of them satisfies a central limit theorem. The application of this to the limiting distribution for a certain class of</p>			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS <input type="checkbox"/>		21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED	
22a. NAME OF RESPONSIBLE INDIVIDUAL Major Brian W. Woodruff		22b. TELEPHONE NUMBER (Include Area Code) (202) 767-5026	22c. OFFICE SYMBOL AFOSR/NM

DD FORM 1473, 83 APR


EDITION OF 1 JAN 73 IS OBSOLETE.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

# CENTER FOR STOCHASTIC PROCESSES

Department of Statistics  
University of North Carolina  
Chapel Hill, North Carolina




## LIMITING DISTRIBUTIONS OF NON-LINEAR VECTOR FUNCTIONS OF STATIONARY GAUSSIAN PROCESSES

by

Hwai-Chung Ho

and

Tze-Chien Sun



Technical Report No. 228

March 1988

145. G. Kalitjanpur and V. Perez-Abreu, Stochastic evolution equations with values on the dual of a countably Hilbert nuclear space, July 86. *Appl. Math. Optimization*, to appear.
146. B.G. Nguyen, Fourier transform of the percolation free energy, July 86. *Probab. Theor. Rel. Fields*, to appear.
147. A.M. Hasofer, Distribution of the maximum of a Gaussian process by Monte Carlo, July 86. *J. Sound Vibration*, 112, 1987, 283-293.
148. T. Norberg, On the existence of probability measures on continuous semi-lattices, Aug. 86.
149. G. Samorodnitsky, Continuity of Gaussian processes, Aug. 86. *Ann. Probability*, to appear.
150. T. Hsing, J. Husler, and M.R. Leadbetter, Limits for exceedance point processes, Sept. 86. *Prob. Theory and Related Fields*, to appear.
151. S. Cambanis, Random filters which preserve the stability of random inputs, Sept. 86. *Adv. Appl. Probability*, 1987, to appear.
152. O. Kallenberg, On the theory of conditioning in point processes, Sept. 86. *Proceedings for the First World Congress of the Bernoulli Society*, Tashkent, 1986.
153. G. Samorodnitsky, Local moduli of continuity for some classes of Gaussian processes, Sept. 86.
154. V. Mandrekar, On the validity of Beurling theorems in polydiscs, Sept. 86.
155. R.F. Serfozo, Extreme values of queue lengths in  $M/G/1$  and  $GI/M/1$  systems, Sept. 86.
156. B.G. Nguyen, Gap exponents for percolation processes with triangle condition, Sept. 86. *J. Statist. Physics*, 49, 1987, 235-243.
157. G. Kalitjanpur and R. Wolpert, Weak convergence of stochastic neuronal models, Oct. 86. *Stochastic Methods in Biology*, M. Kimura et al., eds., Lecture Notes in Biomathematics, 70, Springer, 1987, 116-145.
158. G. Kalitjanpur, Stochastic differential equations in duals of nuclear spaces with some applications, Oct. 86. *Inst. of Math. & Its Applications*, 1986.
159. G. Kalitjanpur and R.L. Karandikar, The filtering problem for infinite dimensional stochastic processes, Jan. 87. *Proc. Workshop on Stochastic Differential Systems, Stochastic Control Theory & Applications*, Springer, to appear.
160. V. Perez-Abreu, Multiple stochastic integrals and nonlinear functionals of a nuclear space valued Wiener process, Oct. 86. *Appl. Math. Optimization*, 16, 1987, 227-245.
161. R.L. Karandikar, On the Feynman-Kac formula and its applications to filtering theory, Oct. 86. *Appl. Math. Optimization*, to appear.
162. R.L. Taylor and T.-C. Hu, Strong laws of large numbers for arrays of rowwise independent random elements, Nov. 86. *Intern. J. Math. & Math. Sci.*, to appear.
163. M. O'Sullivan and T.R. Fleming, Statistics for the two-sample survival analysis problem based on product limit estimators of the survival functions, Nov. 86.
164. F. Avram, On bilinear forms in Gaussian random variables, Toeplitz matrices and Parseval's relation, Nov. 86.
165. D.B.H. Cline, Joint stable attraction of two sums of products, Nov. 86. *J. Multivariate Anal.*, to appear.
166. R.J. Wilson, Model fields in crossing theory—a weak convergence perspective, Nov. 86.
167. D.B.H. Cline, Consistency for least squares regression estimators with infinite variance data, Dec. 86.
168. L.L. Campbell, Phase distribution in a digital frequency modulation receiver, Nov. 86.
169. B.G. Nguyen, Typical cluster size for 2-dim percolation processes, Dec. 86. *J. Statist. Physics*, to appear.
170. H. Odaira, Freidlin-Wentzell type estimates for a class of self-similar processes represented by multiple Wiener integrals, Dec. 86.
171. J. Nolan, Local properties of index- $\beta$  stable fields, Dec. 86. *Ann. Probability*, to appear.
172. R. Menich and R.F. Serfozo, Optimality of shortest queue routing for dependent service stations, Dec. 86.
173. F. Avram and M.S. Taqqu, Probability bounds for  $M$ -Skorohod oscillations, Dec. 86.
174. F. Moricz and R.L. Taylor, Strong laws of large numbers for arrays of orthogonal random variables, Dec. 86.
175. G. Kalitjanpur and V. Perez-Abreu, Stochastic evolution driven by nuclear space valued martingales, Apr. 87.
176. E. Merzbach, Point processes in the plane, Feb. 87.
177. Y. Kasahara, M. Maejima and V. Vervaat, Log fractional stable processes, March 87.
178. G. Kalitjanpur, A.G. Niasse and H. N'kasi, On the prediction theory of two parameter stationary random fields, March 87.
179. R. Brigola, Remark on the multiple Wiener integral, Mar. 87.
180. R. Brigola, Stochastic filtering solutions for ill-posed linear problems and their extension to measurable transformations, Mar. 87.
181. G. Samorodnitsky, Maximal of symmetric stable processes, Mar. 87.
182. H.L. Hurd, Representation of harmonizable periodically correlated processes and their covariance, Apr. 87.
183. H.L. Hurd, Nonparametric time series analysis for periodically correlated processes, Apr. 87.
184. T. Mori and H. Odaira, Freidlin-Wentzell estimates and the law of the iterated logarithm for a class of stochastic processes related to symmetric statistics, May 87.

# LIMITING DISTRIBUTIONS OF NON-LINEAR VECTOR FUNCTIONS OF STATIONARY GAUSSIAN PROCESSES



Hwai-Chung Ho<sup>1</sup>  
Institute of Statistics  
Academia Sinica  
Taipei, Taiwan  
and  
Center for Stochastic Processes  
Department of Statistics  
University of North Carolina  
Chapel Hill, NC 27599-3260

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input checked="" type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
<b>A-1</b>	

and  
Tze-Chien Sun  
Department of Mathematics  
Wayne State University  
Detroit, MI 48202

Abstract: Given a stationary Gaussian vector process  $(X_m, Y_m)$ ,  $m \in \mathbb{Z}$ , and two real functions  $H(x)$  and  $K(x)$  we define  $Z_H^n = A_n^{-1} \sum_{m=0}^{n-1} H(X_m)$  and  $Z_K^n = B_n^{-1} \sum_{m=0}^{n-1} K(Y_m)$ , where  $A_n$  and  $B_n$  are some appropriate constants. The joint limiting distribution of  $(Z_H^n, Z_K^n)$  is investigated. It is shown that  $Z_H^n$  and  $Z_K^n$  are asymptotically independent when one of them satisfies a central limit theorem. The application of this to the limiting distribution for a certain class of non-linear infinite-coordinated functions of a Gaussian process is also discussed.

AMS 1980 subject classification: Primary 60F05, secondary 60G15.

Keywords and phrases: central limit theorem; non-central limit theorem; long-range dependence; stationary Gaussian vector processes.

<sup>1</sup>Research supported by a scholarship received from the National Science Council, ROC, and completed during a visit to the Center for Stochastic Processes, Department of Statistics, University of North Carolina at Chapel Hill.

Let  $(X_m, Y_m)$ ,  $m \in \mathbb{Z}$ , be a sequence of stationary Gaussian vectors. We assume that  $EX_m = EY_m = 0$ ,  $EX_m^2 = EY_m^2 = 1$ .

$$r_1(m) = EX_0 X_m \sim |m|^{-\beta_1},$$

$$r_2(m) = EY_0 Y_m \sim |m|^{-\beta_2},$$

as  $|m| \rightarrow \infty$ , and

$$r_3(m) = EX_0 Y_m \sim m^{-\beta_3},$$

$$r_3(-m) = EY_0 X_m \sim m^{-\beta_4},$$

as  $m \rightarrow \infty$ , where  $\beta_1, \beta_2, \beta_3$  and  $\beta_4 > 0$ . With their correlation functions assumed as above  $\{X_m\}$  and  $\{Y_m\}$  are usually called processes with long-range dependence if  $\beta_1, \beta_2 < 1$ . Let  $G_1(x)$  and  $G_2(x)$  be the spectral distributions of  $\{X_m\}$  and  $\{Y_m\}$ , and let  $Z_{G_1}$  and  $Z_{G_2}$  be their corresponding random measures. Since  $\{(X_m, Y_m)\}$  is stationary there always exists a complex-valued function  $G_3(x)$  such that

$$r_3(m) = \int e^{-imx} dG_3(x), \quad \forall m \in \mathbb{Z}.$$

Since the matrix

$$\begin{bmatrix} G_1(dx) & G_3(dx) \\ \overline{G_3}(dx) & G_2(dx) \end{bmatrix}$$

is positive definite, it follows that

$$(1) \quad |G_3(dx)|^2 \leq G_1(dx)G_2(dx).$$

Given two functions  $H(x)$  and  $K(x)$ , satisfying  $EH(X_m) = EK(Y_m) = 0$ ,  $EH^2(X_m) < \infty$

and  $EK^2(Y_m) < \infty$ , and having their Hermite expansions as follows:

$$H(x) = \sum_{j=\nu_1}^{\infty} c_j H_j(x) \quad \text{and} \quad K(x) = \sum_{j=\nu_2}^{\infty} d_j H_j(x),$$

we define

$$Z_H^n = A_n^{-1} \sum_{m=0}^{n-1} H(X_m) \quad \text{and} \quad Z_K^n = B_n^{-1} \sum_{m=0}^{n-1} K(Y_m).$$

It has been proved that with proper choice of the norming factors  $A_n$  and  $B_n$ , then as  $n \rightarrow \infty$ ,  $Z_H^n$  and  $Z_K^n$  have limiting distributions. When  $\nu_i \beta_i < 1$ ,  $i=1,2$ , the limiting distribution is non-Gaussian (unless  $\nu_i=1$ ) and can be represented by multiple Wiener integrals ([2], [8]). When the limit law is non-Gaussian, it is usually said that a non-central limit theorem (or NCLT) is satisfied. On the other hand a central limit theorem (or CLT, i.e. the norming factor is  $n^{1/2}$  and the limit law is Gaussian) will hold if  $\nu_i \beta_i > 1$  [1]. The purpose of this paper is to study the joint limiting distribution of  $(Z_H^n, Z_K^n)$ , when, in particular, one component satisfies a CLT. An incomplete attempt at solving the same problem had been made by Hsiao [4].

The reason we look into this problem is the following: Consider the following square-integrable function  $L(\cdot)$  (possibly infinite - coordinated) of a stationary Gaussian process, defined by its Wiener-Ito expansion

$$L = I_{k_1}(f_1) + I_{k_2}(f_2), \quad 1 \leq k_1 < k_2,$$

where  $I_j(f)$  is the  $j$ -fold multiple Wiener integral with kernel  $f$ . Let  $Z_L^n$  be defined as

$$\begin{aligned} Z_L^n &= C_n^{-1} \sum_{m=0}^{n-1} I_{k_1}(U_m \circ f_1) + C_n^{-1} \sum_{m=0}^{n-1} I_{k_2}(U_m \circ f_2) \\ &\equiv Z_{L_1}^n + Z_{L_2}^n \end{aligned}$$



where  $U_m$  is the  $m$ -step shift operator, i.e.  $(U_m \circ f)(x_1, \dots, x_k) = \exp(im(x_1 + \dots + x_k))f(x_1, \dots, x_k)$ . Previous studies on the limit laws of  $Z_L^n$  are mainly directed to the cases where: both  $Z_{L_1}^n$  and  $Z_{L_2}^n$  satisfy either a CLT ([1], [3]) or a NCLT [7]. The case where one of  $Z_{L_1}^n$  and  $Z_{L_2}^n$  satisfies a CLT and the other one satisfies a NCLT is still unclear, and the following two natural questions arise: Will the limit law (if it exists)  $Z_L^\infty$  of  $Z_L^n$  be still equal to the sum of the limit laws  $Z_{L_1}^\infty$  and  $Z_{L_2}^\infty$  of  $Z_{L_1}^n$  and  $Z_{L_2}^n$ ; and what is the relation between  $Z_{L_1}^\infty$  and  $Z_{L_2}^\infty$ . The main result of this paper, stated in the Theorem, provides an answer to these questions. Suppose the underlying stationary Gaussian process for  $Z_L^n$  exhibits long-range dependence. For a certain class of functions  $L(\cdot)$ , by making use of the formula for the change variables ([5], p. 32) on the kernels  $f_1$  and  $f_2$ , we may obtain

$$(Z_{L_1}^n, Z_{L_2}^n) \stackrel{\Delta}{=} (C_n^{-1} \sum_{m=0}^{n-1} H_{k_1}(Y'_m), C_n^{-1} \sum_{m=0}^{n-1} H_{k_2}(X'_m)),$$

for some sequence of stationary Gaussian vectors  $(X'_m, Y'_m)$ ,  $m \in \mathbb{Z}$ . " $\stackrel{\Delta}{=}$ " means equal in distribution. Assume  $C_n = n^{1/2}$ . Suppose  $Z_{L_1}^n$  and  $Z_{L_2}^n$  satisfy a CLT and a NCLT respectively. If, furthermore, the conditions in the Theorem are met by  $\{(X'_m, Y'_m)\}$ , then as a result of the Theorem, it follows that

$$Z_{L_1}^\infty \perp Z_{L_2}^\infty \quad \text{and} \quad Z_L^\infty \stackrel{\Delta}{=} Z_{L_1}^\infty + Z_{L_2}^\infty$$

(" $\perp$ " means independent), i.e. the distribution function  $\tilde{L}(x)$  of  $Z_L^\infty$  can be written as

$$\tilde{L}(x) = \int F(x-y) d\Phi(y/\sigma), \quad \sigma > 0.$$

for some distribution function  $F(y)$  and a standard Gaussian distribution  $\Phi(y)$ . It should be pointed out that a NCLT with norming factor  $n^{1/2}$ , such as for  $Z_{L_2}^n$ , is shown possible in [6]. A more detailed study of the situation where a CLT and a NCLT jointly occur will appear in a subsequent paper by the authors. Now we formulate our main result:

Theorem. Assume  $v_1\beta_1 < 1 < v_2\beta_2$ . When  $v_2 = 1$  we also assume

$$\beta \equiv \beta_3 \wedge \beta_4 > \frac{1 + \beta_1}{2}.$$

Then with  $A_n = n^{1-v_1\beta_1/2}$  and  $B_n = n^{1/2}$  the limiting distributions  $Z_H^*$  and  $Z_K^*$  of  $Z_H^n$  and  $Z_K^n$  are independent.

Note that  $Z_K^*$  is Gaussian by [1].

Throughout the rest of the paper we always assume  $v_1\beta_1 < 1$  and  $v_2\beta_2 > 1$ . Later, in proving the Theorem, we shall only deal with the very special case where  $H(x)$  and  $K(x)$  have the following one-term expansion

$$H(x) = H_{v_1}(x) \quad \text{and} \quad K(x) = H_{v_2}(x).$$

The reduction of  $H(x)$  to its first term is justified because when  $v_1\beta_1 < 1$  only the first term is relevant to the distribution  $Z_H^*$  [8]. In [1] it is made clear that when  $v_2\beta_2 > 1$  we need only to consider the  $K(x)$  with finite expansion to prove the central limit theorem. Though we prove the Theorem only for the  $K(x)$  with one-term expansion, the arguments in the proof can be easily extended to the finite expansion case.

The major tool we use to prove the Theorem is the so-called "diagram formula" [5] on how to compute the expectation of a product of Hermite polynomials of standard Gaussian random variables. Prior to giving the

statement of the formula, we need some notations and definitions. Let a given set of  $(\ell_1 + \dots + \ell_p)$  vertices be arranged into  $p$  levels such that the  $i$ -th level has  $\ell_i$  vertices. A graph  $G$  is called a diagram of order  $(\ell_1, \dots, \ell_p)$  if (1) each vertex is of degree one and (2) edges may pass only between different levels. By a regular diagram we mean a diagram whose edges do not pass between levels in different pairs. For each edge  $w \in G$  connecting the  $i$ -th and  $j$ -th level,  $i < j$ , define  $d_1(w) = i$  and  $d_2(w) = j$ .

**Lemma 1.** (Diagram Formula) Let  $(W_1, \dots, W_p)$  be a Gaussian vector with  $EW_1 = 0$ ,  $EW_1^2 = 1$ , and  $EW_1 W_j = r(1, j)$ . Then for the Hermite polynomials  $H_{\ell_1}(x), \dots, H_{\ell_p}(x)$ , we have

$$E \prod_{i=1}^p H_{\ell_i}(W_i) = \sum_{G \in \mathcal{G}} \prod_{w \in G} r(d_1(w), d_2(w)),$$

where the sum runs through all the diagrams  $G$  of order  $(\ell_1, \dots, \ell_p)$ .

The following lemma is well-known and can be easily derived from Lemma 1.

**Lemma 2.** Given two r.v.'s  $Z$  and  $W$  with  $EZ = EW = 0$ ,  $EZ^2 = \sigma_1^2$  and  $EW^2 = \sigma_2^2$ , then  $Z$  and  $W$  are independent Gaussian r.v.'s if and only if for all  $\ell, m$ ,

$$EZ^\ell W^m = \begin{cases} \frac{\ell! m!}{2^{\ell/2+m/2} (\ell/2)! (m/2)!} \sigma_1^\ell \sigma_2^m & \text{if } \ell \text{ and } m \text{ are even} \\ 0 & \text{otherwise.} \end{cases}$$

In the following " $\Delta$ " always denotes bounded Borel sets in  $\mathbb{R}$ . Then because  $r_3(n) = \int e^{-inx} dG_3(x)$ , we have

$$(2) \quad E \int f(x) Z_{G_1}(dx) \int g(x) Z_{G_2}(dx) = \int f(x) \bar{g}(x) dG_3(x)$$

for  $f \in L^2(G_1)$  and  $g \in L^2(G_2)$ . By Proposition 1 of [2] or similar arguments it

can be proved that there exist  $G_1^*(x)$  and  $G_2^*(x)$  such that

$$(3.1) \quad n^{\beta_1} G_1\left(\frac{dx}{n}\right) \xrightarrow{\text{weakly}} G_1^*(dx)$$

and

$$(3.2) \quad n^{\beta_2^*} (\log n)^{-\delta(\beta_2)} G_2\left(\frac{dx}{n}\right) \xrightarrow{\text{weakly}} G_2^*(dx)$$

as  $n \rightarrow \infty$ , where  $\beta_2^* = \beta_2 \wedge 1$  and  $\delta(x) = 1$  if  $x=1$ , and  $= 0$  if  $x \neq 1$ .

We shall need the following lemma to prove the Theorem. Recall that  $\beta = \beta_3 \wedge \beta_4$ .

**Lemma 3.** Assume  $\beta \leq 1$ . There exists a function  $G_3^*(x)$  of locally bounded variation such that for each bounded Borel set  $\Delta$ ,

$$(4) \quad \lim_{m \rightarrow \infty} m^\beta (\log m)^{-\delta(\beta)} G_3\left(\frac{\Delta}{m}\right) = G_3^*(\Delta).$$

Moreover  $G_3^*$  satisfies

$$G_3^*([0, y]) = \overline{G_3([-y, 0])} = y^\beta D,$$

where  $D$  is some complex constant.

**Proof:** It is sufficient to show that (4) holds for  $\Delta = [0, y]$  or  $[-y, 0]$ .

Define

$$F_n(x) = \frac{1}{2\pi} \sum_{|s| \leq n} r_3(x) \cdot \int_{-\pi}^x e^{isy} dy$$

for  $x \in [-\pi, \pi]$ . Since each term in the above sum is bounded by  $C \cdot |s|^{-\beta-1}$  for some constant  $C$ ,  $F_n(x)$  converges to  $G_3(x)$  for all  $x$ . i.e.

$$G_3(x) = \lim_{n \rightarrow \infty} F_n(x) = \frac{1}{2\pi} \sum_{s=-\infty}^{\infty} r_3(s) \frac{e^{isx} - e^{-is\pi}}{is}.$$

Let  $\Delta = [0, y]$ . Define

$$S_{m,y} = m^\beta (\log m)^{-\delta(\beta)} [G_3(\frac{\Delta}{m}) + \overline{G_3(\frac{\Delta}{m})}].$$

By  $\overline{G_3(\Delta)} = G_3(-\Delta)$ ,  $S_{m,y}$  is equal to

$$S_{m,y} = \frac{1}{\pi} (\log m)^{-\delta(\beta)} \sum_{s=-\infty}^{\infty} (r_3(s)m^\beta) \frac{\sin \frac{s}{m} y}{s},$$

which as  $m \rightarrow \infty$  tends to

$$(4.1) \quad \lim_{m \rightarrow \infty} S_{m,y} = \begin{cases} y^\beta R_\beta & \text{if } \beta_3 \neq \beta_4, \\ y^\beta 2R_\beta & \text{if } \beta_3 = \beta_4, \end{cases}$$

where  $R_\beta = \frac{1}{\pi} \lim_{\epsilon \rightarrow 0} (-\log \epsilon)^{-\delta(\beta)} \int_\epsilon^\infty x^{-\beta-1} \sin x \, dx$ . Similarly if we define

$$\begin{aligned} C_{m,y} &= m^\beta (\log m)^{-\delta(\beta)} [G_3(\frac{\Delta}{m}) - \overline{G_3(\frac{\Delta}{m})}] \\ &= i \frac{1}{\pi} (\log m)^{-\delta(\beta)} \sum_{s=-\infty}^{\infty} (r_3(s)m^\beta) \frac{(1 - \cos \frac{s}{m} y)}{s} \end{aligned}$$

then we obtain

$$(4.2) \quad \lim_{m \rightarrow \infty} C_{m,y} = \begin{cases} -iy^\beta I_\beta & \text{if } \beta_3 < \beta_4 \\ iy^\beta I_\beta & \text{if } \beta_4 < \beta_3 \\ 0 & \text{if } \beta_3 = \beta_4 \end{cases}$$

where  $I_\beta = \frac{1}{\pi} \lim_{\epsilon \rightarrow 0} (-\log \epsilon)^{-\delta(\beta)} \int_\epsilon^\infty x^{-\beta-1} (1 - \cos x) dx$ . (4.1) and (4.2) imply

$$\lim_{m \rightarrow \infty} m^\beta (\log m)^{-\delta(\beta)} G_3\left(\frac{[0,y]}{m}\right) = y^\beta D \equiv G_3^*([0,y]).$$

where

$$D \equiv \begin{cases} y^{\beta} R_{\beta} & \text{if } \beta_3 = \beta_4 \\ y^{\beta} (R_{\beta} - iI_{\beta})/2 & \text{if } \beta_3 < \beta_4 \\ y^{\beta} (R_{\beta} + iI_{\beta})/2 & \text{if } \beta_4 < \beta_3 \end{cases}.$$

Since the property  $\overline{G_3(\Delta)} = G_3(-\Delta)$  is preserved by passing to the limit  $G_3^*$ , we have

$$G_3^*([-y, 0]) = y^{\beta} \overline{D}.$$

The proof is completed. □

Assume  $\beta \leq 1$  again and observe that

$$\begin{aligned} & m^{\beta} (\log m)^{-\delta(\beta)} |G_3(\frac{\Delta}{m})| \\ & \leq m^{\beta - (\beta_1 + \beta_2^*)/2} (\log m)^{\delta(\beta_2) - \delta(\beta)} [m^{\beta_1} G_1(\frac{\Delta}{m}) m^{\beta_2^*} (\log m)^{-\delta(\beta_2)} G_2(\frac{\Delta}{m})]^{1/2}. \end{aligned}$$

Then we have an immediate corollary from (3.1), (3.2) and (4):

$$(4.3) \quad \beta \geq (\beta_1 + \beta_2^*)/2.$$

When  $\beta > 1$ , then  $G_3(dx)$  is absolutely continuous and its density is continuous.

Let  $G_3(dx) = f(x)dx$ . Then

$$(4.4) \quad \lim_{m \rightarrow \infty} m G_3(\frac{\Delta}{m}) = \lambda(\Delta) f(0),$$

where  $\lambda$  is the Lebesgue measure. (4.3) is clearly satisfied for  $\beta > 1$ .

By (3.1),

$$n^{1-\beta_1/2} Z_{G_1}(\frac{\Delta}{n}) \xrightarrow{d} Z_{G_1^*}(\Delta)$$

as  $n \rightarrow \infty$  [2], where  $Z_{G_1^*}$  is the random measure induced by  $G_1^*(dx)$ . Since the

distribution of  $Z_H^*$  can be represented by the  $v_1$ -fold Wiener integral, we need to show that for disjoint  $\Delta_i$ 's,  $i=1,2,\dots,v_1$ ,

$$(Z_{G_1}^*(\Delta_1), \dots, Z_{G_1}^*(\Delta_{v_1})) \perp Z_k^*,$$

which is in fact equivalent to showing for each  $\Delta$ ,

$$(5) \quad Z_{G_1}^*(\Delta) \perp Z_k^*.$$

It is not difficult to see that (5) is equivalent to

$$(5.1) \quad W(\Delta) = \int_{\Delta} \frac{e^{ix} - 1}{ix} Z_{G_1}^*(dx) \perp Z_k^*.$$

It is mere technical convenience that leads us to replace  $Z_{G_1}^*(\Delta)$  by  $W(\Delta)$ .

Define

$$(5.2) \quad K_n(x) \equiv \frac{e^{ix} - 1}{(e^{ix/n} - 1)n} = \frac{1}{n} \sum_{j=0}^{n-1} e^{ijx/n}$$

and

$$\begin{aligned} W_n(\Delta) &\equiv n^{-(1-\beta_1/2)} \sum_{j=0}^{n-1} \int_{\Delta} e^{ijx} Z_{G_1}^*(dx) \\ &= \int_{\Delta} K_n(x) n^{-(1-\beta_1/2)} Z_{G_1}^*\left(\frac{dx}{n}\right). \end{aligned}$$

(3.1) and the fact that  $K_n(x)$  converges to  $(e^{ix}-1)/ix$  uniformly on every bounded set imply

$$(6) \quad W_n(\Delta) \xrightarrow{d} W(\Delta)$$

as  $n \rightarrow \infty$  [2]. If we can show

$$\begin{aligned}
 & \lim_{n \rightarrow \infty} E(W_n(\Delta))^{\ell} (Z_k^n)^m \\
 (7) \quad & = \begin{cases} \frac{\ell! m!}{2^{\ell/2 + m/2} (\ell/2)! (m/2)!} \sigma_1^{\ell} \sigma_2^m & \text{if } \ell \text{ and } m \text{ are even.} \\ 0 & \text{otherwise} \end{cases}
 \end{aligned}$$

then by (6) and Lemma 2, (5.1) follows. Notice that  $EW^2(\Delta) \equiv \sigma_1^2$  and  $E(Z_k^*)^2 = \sigma_2^2$ .

Proof of Theorem.

Given a fixed setting of vertices  $V = (1, \dots, 1, v_2, \dots, v_2)$  having as follows its configuration

$$\begin{array}{c}
 \ell \quad \left\{ \begin{array}{c} 0 \\ 0 \\ \vdots \\ 0 \end{array} \right. \\
 \\
 m \quad \left\{ \begin{array}{cccc} 0 & 0 & \dots & 0 \\ 0 & 0 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \dots & 0 \end{array} \right. \\
 \\
 v_2
 \end{array}$$

Define  $\Gamma$  = the set of all regular diagrams of order  $V$  and  $\Gamma^c$  the complement of  $\Gamma$ , i.e., the set of all non-regular diagrams of order  $V$ . Any subgraph of a diagram is called a subdiagram if it is itself a diagram and is the union of levels and the edges on the levels. Any diagram  $G \in \Gamma^c$  can be partitioned into three disjoint subdiagrams  $V_{G,1}$ ,  $V_{G,2}$  and  $V_{G,3}$ , which are defined as follows.



$V_{G,1}$  = the maximal subdiagram of  $G$  which is regular within itself, and all its edges satisfy  $1 \leq d_1(w) < d_2(w) \leq \ell$  or  $\ell + 1 \leq d_1(w) < d_2(w) \leq \ell + m$ .

$V_{G,2}$  = the maximal subdiagram of  $G - G_{G,1}$  whose edges satisfy  $\ell + 1 \leq d_1(w) < d_2(w) \leq \ell + m$ .

$V_{G,3} = G - (V_{G,1} \cup V_{G,2})$ .

For each subdiagram  $V_{G,i}$  of  $G$ ,  $i=1,2,3$ , define

$V_{G,i}^* = \{j \mid \text{the } j\text{-th level of } V \text{ is in } V_{G,i}\}$ .

$V_{G,i}^*(1) = \{j \in V_{G,i}^* \mid 1 \leq j \leq \ell\}$ .

$V_{G,i}^*(2) = \{j \in V_{G,i}^* \mid \ell + 1 \leq j \leq \ell + m\}$ .

In the following  $E(G)$  denotes the set of all edges contained in the diagram  $G$ .

Use Lemma 2 (Diagram Formula)

$$\begin{aligned}
 E(W_n(\Delta))^{\ell} (Z_k^n)^m &= \sum_{G \in \Gamma} [E(W_n(\Delta))^2]^{\ell/2} [E(Z_k^n)^2]^{m/2} \\
 &\quad + \sum_{G \in \Gamma^c} \left[ \prod_{w \in E(V_{G,1})} E(W_n(\Delta))^2 \prod_{w \in E(V_{G,1})} E(Z_k^n)^2 \right] \\
 &\quad \times [n^{-|V_{G,2}^*(2)|/2} \sum_{0 \leq p_1 \leq n-1} \prod_{w \in E(V_{G,2})} r_2(p_{d_1(w)} - p_{d_2(w)})] \\
 &\quad \times [n^{-|V_{G,3}^*(2)|/2} \sum_{0 \leq p_1 \leq n-1} \prod_{w \in E(V_{G,3})} r_2(p_{d_1(w)} - p_{d_2(w)})] \\
 &\quad \times \prod_{\substack{w \in E(V_{G,3}) \\ d_1(w) \in V_{G,3}^*(1)}} n^{-(1-\beta_1/2)} E\left(\int_{\frac{\Delta}{n}} \exp(ip_{d_1(w)}x) Z_{G_1}(dx) \int \exp(ip_{d_2(w)}x) Z_{G_2}(dx)\right)
 \end{aligned}$$

$$(8) \quad \equiv \sum_{\Gamma}^n + \sum_{G \in \Gamma^c} A_1^n \times A_2^n \times A_3^n.$$

Since  $\sum_{\Gamma}^n$  converges to the right hand side of (7), it is sufficient to show that for fixed  $G \in \Gamma^c$  the second term of (8) vanishes.

$$(9) \quad \lim_{n \rightarrow \infty} A_1^n \times A_2^n \times A_3^n = 0.$$

Recall the definition of  $W_n(\Delta)$ . We have

$$(10) \quad EW_n^2(\Delta) \rightarrow \int_0 \left| \frac{e^{ix} - 1}{ix} \right|^2 dG_1^*(x) = EW^2(\Delta).$$

as  $n \rightarrow \infty$ . (10) and the central limit theorem for  $Z_k^n$  imply

$$(11) \quad \lim_{n \rightarrow \infty} A_1^n = (EW^2(\Delta))^{|V_{G,1}^*(1)|/2} (\sigma_2^2)^{|V_{G,1}^*(2)|/2}.$$

Using (2) and (5.2), we can rewrite

$$(12) \quad A_3^n = n^{-|V_{G,3}^*(2)|/2} \sum_{\substack{0 \leq p_1 \leq n-1 \\ i \in V_{G,3}^*(2)}} \prod_{\substack{w \in E(V_{G,3}) \\ d_1(w) \in V_{G,3}^*(2)}} r_2(p_{d_1(w)} - p_{d_2(w)}) \cdot \\ \cdot \prod_{\substack{e \in E(V_{G,3}) \\ d_1(e) \in V_{G,3}^*(1)}} \left[ \sum_{0 \leq p_{d_1(e)} \leq n-1} n^{-(1-\beta_1/2)} \int_{\Delta} e^{i \left( \frac{p_{d_1(e)} - p_{d_2(e)}}{n} \right) x} dG_3\left(\frac{x}{n}\right) \right].$$

Fix  $p_1, i \in V_{G,3}^*(2)$ , and  $e \in E(V_{G,3})$  with  $d_1(e) \leq \ell$ . we obtain as a result of (4) in Lemma 3 (or (4.4) if  $\beta > 1$ ) an asymptotic bound for the second summation (denoted by  $\Sigma_n^*$ ) in (12). In the following  $\alpha \equiv \beta \wedge 1$ .

$$\begin{aligned}
\Sigma_n^* &= n^{(\beta_1 - 2\alpha)/2} (\log n)^{\delta(\beta)} \cdot \int_{\Delta} \left[ \sum_{0 \leq p_{d_1(e)} \leq n-1} \exp(ip_{d_1(e)} x/n) \cdot \frac{1}{n} \right] \cdot \\
&\quad \cdot \exp(-ip_{d_2(e)} x/n) \cdot n^{\alpha} (\log n)^{-\delta(\beta)} dG_3\left(\frac{x}{n}\right). \\
&= O(n^{(\beta_1 - 2\alpha)/2} (\log n)^{\delta(\beta)} \int_{\Delta} \left| \frac{e^{ix} - 1}{ix} \right| |dG_3^*(x)|).
\end{aligned}$$

If  $\beta \leq 1$ , then

$$(13) \quad \beta_1 - 2\alpha \leq \beta_1 - 2\beta \leq -\beta_2^* < 0,$$

where we make use of the fact that  $\beta \geq (\beta_1 + \beta_2^*)/2$  derived right after Lemma 3.

If  $\beta > 1$ , clearly (13) still holds. By (13) it follows that

$$(13.1) \quad \Sigma_n^* = o(1).$$

Define

$k(i)$  = the number of edges  $w$  satisfying  $d_1(w) = i$ .

and

$g(i)$  = the number of vertices in the  $i$ -th level not connecting any of the first  $\ell$  levels.

We firstly assume that  $v_2 > 1$ . By the similar argument employed to prove Proposition in [1], we can develop the following facts:

$$(14) \quad \lim_{n \rightarrow \infty} A_2^n = 0$$

if  $V_{G,2}$  is nonempty. And secondly an asymptotic bound as (17) can be obtained for the first summation in (12) if  $V_{G,3} \neq \emptyset$ .

$$\begin{aligned}
 (15) \quad \alpha_n &\equiv \sum_{0 \leq p_1 \leq n-1} \prod_{\substack{w \in E(V_{G,2}) \\ d_1(w) \in V_{G,3}^*(2)}} |r_2(p_{d_1(w)} - p_{d_2(w)})| \\
 &= O(n^{|V_{G,3}^*(2)| - \sum_{i \in V_{G,3}^*(2)} \frac{k(i)}{g(i)}})
 \end{aligned}$$

Note that the  $\alpha_n$  given above is well-defined because it is assumed that  $v_2 > 1$ .

As shown in (2.20) in [1] we have the following inequality

$$(16) \quad \sum_{i \in V_{G,3}^*(2)} \frac{k(i)}{g(i)} \geq \frac{1}{2} |V_{G,3}^*(2)|.$$

(15) and (16) imply that

$$(17) \quad \alpha_n = O(n^{|V_{G,3}^*(2)|/2}).$$

Then (12), (13.1) and (17) imply

$$A_3^n = (o(1))^{|V_{G,3}^*(1)|}.$$

Hence if  $|V_{G,3}^*(1)| > 0$  (because  $V_{G,3} \neq \emptyset$ ), then

$$(18) \quad \lim_{n \rightarrow \infty} A_3^n = 0.$$

For any non-regular diagram  $G \in \Gamma^C$  if  $v_2 > 1$  then its subdiagrams  $V_{G,2}$  and  $V_{G,3}$  can not be empty at the same time, that is either (14) or (18) must hold.

Hence (9) is true.

When  $v_2 = 1$ , then  $V_{G,2}$  is empty, i.e.  $A_2^n$  is absent. Thus in order to assure (9) we have to show (18). Also note that when  $v_2 = 1$  the first product in (12) no longer exists. Rewrite  $A_3^n$  given in (12) in a more simplified form and apply to it the result of Lemma 3. We have

$$\begin{aligned}
A_3^n &= \prod_{w \in E(V_{G,3})} n^{(\beta_1+1)/2-\alpha} (\log n)^{\delta(\beta)} \\
&\cdot \int_{\Delta} \left[ \sum_{0 \leq p_{d_1(w)} \leq n-1} \exp(ip_{d_1(w)} x/n) \cdot \frac{1}{n} \right] \cdot \\
&\cdot \left[ \sum_{0 \leq p_{d_2(w)} \leq n-1} \exp(-ip_{d_2(w)} x/n) \cdot \frac{1}{n} \right] n^{\alpha} (\log n)^{-\delta(\beta)} dG_3\left(\frac{x}{n}\right) \\
&= O(n^{(\beta_1+1)/2-\alpha} (\log n)^{\delta(\beta)} \int_{\Delta} \left| \frac{e^{ix} - 1}{ix} \right|^2 |dG_3^*(x)|).
\end{aligned}$$

By the assumption of the Theorem, when  $v_2 = 1$ ,

$$\begin{aligned}
\frac{\beta_1 + 1}{2} < \beta &\Rightarrow \frac{\beta_1 + 1}{2} - \alpha < 0, \\
&\Rightarrow \lim_{n \rightarrow \infty} A_3^n = 0.
\end{aligned}$$

The proof is completed. □

## REFERENCES

- [1] Breuer, P., and Major, P. (1983). Central limit theorems for non-linear functionals of Gaussian fields, J. Multivariate Anal., 13, 425-441.
- [2] Dobrushin, R.L., and Major, P. (1979). Non-Central Limit theorems for non-linear functionals of Gaussian fields, Z. Wahrsch. verw. Gebiete 50, 27-52.
- [3] Ho, H.C., and Sun, T.C. (1987). A central limit theorem for non-instantaneous filters of a stationary Gaussian Process, J. Multivariate Anal., 22, 144-155.
- [4] Hsiao, C.T. (1983). Central limit theorem for stationary random vectors, unpublished note.
- [5] Major, P. (1981). Multiple Wiener Integrals, Lecture Notes in Math. 849, Springer-Verlag, New York/Berlin.
- [6] Major, P. (1981). Limit theorems for non-linear functionals of Gaussian sequences, Z. Wahrsch. verw. Gebiete 57, 129-158.
- [7] Rosenblatt, M. (1987). Remarks on limit theorems for non-linear functionals of Gaussian sequences, Probab. Th. Rel. Fields, 75, 1-10.
- [8] Taqqu, M. (1979). Convergence of iterated processes of arbitrary Hermit rank, Z. Wahrsch. verw. Gebiete 50, 53-83.

185. R.F. Serfozo, Point processes, May 87. Operations Research Handbook on Stochastic Processes, to appear.
186. Z.D. Bai, W.Q. Liang and W. Vervaat, Strong representation of weak convergence, June 87.
187. O. Kallenberg, Decoupling identities and predictable transformations in exchangeability, June, 87.
188. O. Kallenberg, An elementary approach to the Daniell-Kolmogorov theorem and some related results, June 87. Math. Nachr., to appear.
189. G. Samorodnitsky, Extrema of skewed stable processes, June 87.
190. D. Nualart, M. Sanz and M. Zakai, On the relations between increasing functions associated with two-parameter continuous martingales, June 87.
191. F. Avram and M. Taqqu, Weak convergence of sums of moving averages in the  $\alpha$ -stable domain of attraction, June 87.
192. M.R. Leadbetter, Harald Cramér (1893-1985), July 87. ISI Review, to appear.
193. R. LePage, Predicting transforms of stable noise, July 87.
194. R. LePage and B.M. Schreiber, Strategies based on maximizing expected log, July 87.
195. J. Rosinski, Series representations of infinitely divisible random vectors and a generalized shot noise in Banach spaces, July 87.
196. J. Szulga, On hypercontractivity of  $\alpha$ -stable random variables,  $0 < \alpha < 2$ , July 87.
197. I. Kuznezova-Sholpo and S.T. Rachev, Explicit solutions of moment problems I, July 87. Probability and Mathematical Statistics, 10, 1989, to appear.
198. T. Hsing, On the extreme order statistics for a stationary sequence, July 87.
199. T. Hsing, Characterization of certain point processes, Aug. 87. Stochastic Proc. Appl. 26, 1987, 297-316.
200. J.P. Nolan, Continuity of symmetric stable processes, Aug. 87.
201. M. Marques and S. Cambanis, Admissible and singular translates of stable processes, Aug. 87.
202. O. Kallenberg, One-dimensional uniqueness and convergence results for exchangeable processes, Aug. 87.
203. R.J. Adler, S. Cambanis and G. Samorodnitsky, On stable Markov processes, Sept. 87.
204. G. Kallianpur and V. Perez-Abreu, Stochastic evolution equations driven by nuclear space valued martingales, Sept. 87.
205. R.L. Smith, Approximations in extreme value theory, Sept. 87.
206. E. Willemens, Estimation of convolution tails, Sept. 87.
207. J. Rosinski, On path properties of certain infinitely divisible processes, Sept. 87.

208. A.H. Koreziloglu, Computation of filters by sampling and quantization, Sept. 87.
209. J. Bather, Stopping rules and observed significance levels, Sept. 87.
210. S.T. Rachev and J.E. Yukich, Convolution metrics and rates of convergence in the central limit theorem, Sept. 87.
211. M. Fujisaki, Normed Bellman equation with degenerate diffusion coefficients and its applications to differential equations, Oct. 87.
212. C. Simons, Y.C. Yao and X. Wu, Sequential tests for the drift of a Wiener process with a smooth prior, and the heat equation, Oct. 87.
213. R.L. Smith, Extreme value theory for dependent sequences via the Stein-Chen method of Poisson approximation, Oct. 87.
214. C. Houdré, A note on vector bimeasures, Nov. 87.
215. M.R. Leadbetter, On the exceedance random measures for stationary processes, Nov. 87.
216. M. Marques, A study on Lebesgue decomposition of measures induced by stable processes, Nov. 87.
217. M.T. Alpuim, High level exceedances in stationary sequences with extremal index, Dec. 87.
218. R.F. Serfozo, Poisson functionals of Markov processes and queueing networks, Dec. 87.
219. J. Bather, Stopping rules and ordered families of distributions, Dec. 87.
220. S. Cambanis and M. Masjima, Two classes of self-similar stable processes with stationary increments, Jan. 88.
221. H.P. Huckle, G. Kallianpur and R.L. Karandikar, Smoothness properties of the conditional expectation in finitely additive white noise filtering, Jan. 88.
222. I. Mitoma, Weak solution of the Langevin equation on a generalized functional space, Feb. 88.
223. L. de Haan, S.I. Resnick, H. Rootzén and C. de Vries, Extremal behaviour of solutions to a stochastic difference equation with applications to arch-processes, Feb. 88.
224. O. Kallenberg and J. Szulga, Multiple integration with respect to Poisson and Lévy processes, Feb. 88.
225. D.A. Dawson and L.C. Gorostiza, Generalized solutions of a class of nuclear space valued stochastic evolution equations, Feb. 88.
226. C. Samorodnitsky and J. Szulga, An asymptotic evaluation of the tail of a multiple symmetric  $\alpha$ -stable integral, Feb. 88.
227. J.J. Hunter, The computation of stationary distributions of Markov chains through perturbations, Mar. 88.
228. H.C. Ho and T.C. Sun, Limiting distribution of nonlinear vector functions of stationary Gaussian processes, Mar. 88.

END

DATE

FILMED

8-88

DTIC